



Fermi National Accelerator Laboratory

**FERMILAB-Pub-93/323-E
E687**

An Observation of an Excited State of the Λ_c^+ Baryon

**P.L. Frabetti et al
The E687 Collaboration**

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

November 1993

Submitted to *Physical Review Letters*

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

An Observation of an Excited State of the Λ_c^+ Baryon

P. L. Frabetti

Dip. di Fisica dell'Università and INFN - Bologna, I-40126 Bologna, Italy

H. W. K. Cheung, J. P. Cumalat, C. Dallapiccola*, J. F. Ginkel, S. V. Greene, W. E. Johns, M. S. Nehring
University of Colorado, Boulder, CO 80309

J. N. Butler, S. Cihangir, I. Gaines, P. H. Garbincius, L. Garren, S. A. Gourlay, D. J. Harding, P. Kasper,
A. Kreymer, P. Lebrun, S. Shukla
Fermilab, Batavia, IL 60510

S. Bianco, F. L. Fabbri, S. Sarwar, A. Zallo
Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

R. Culbertson, R. W. Gardner, R. Greene, J. Wiss
University of Illinois at Urbana-Champaign, Urbana, IL 61801

G. Alimonti, G. Bellini, B. Caccianiga, L. Cinquini[†], M. Di Corato, M. Giammarchi, P. Inzani, F. Leveraro,
S. Malvezzi[‡], D. Menasce, E. Meroni, L. Moroni, D. Pedrini, L. Perasso, A. Sala, S. Sala, D. Torretta[§], M. Vittono[§]
Dip. di Fisica dell'Università and INFN - Milano, I-20133 Milan, Italy

D. Buchholz, D. Claes, B. Gobbi, B. O'Reilly
Northwestern University, Evanston, IL 60208

J. M. Bishop, N. M. Cason, C. J. Kennedy**, G. N. Kim, T. F. Lin, D. L. Pusejlic, R. C. Ruchti, W. D. Shephard,
J. A. Swiatek, Z. Y. Wu
University of Notre Dame, Notre Dame, IN 46556

V. Arena, G. Boca, C. Castoldi, G. Gianini, S. P. Ratti, C. Riccardi, P. Vitulo
Dip. di Fisica Nucleare e Teorica dell'Università and INFN - Pavia, I-27100 Pavia, Italy.

A. Lopez
University of Puerto Rico at Mayaguez, Puerto Rico

G. P. Grim, V. S. Paolone, P. M. Yager
University of California-Davis, Davis, CA 95616

J. R. Wilson
University of South Carolina, Columbia, SC 29208

P. D. Sheldon
Vanderbilt University, Nashville, TN 37235

F. Davenport
University of North Carolina-Asheville, Asheville, NC 28804

J. F. Filasetta
Northern Kentucky University, Highland Heights, KY 41076

G.R. Blackett, M. Pisharody, T. Handler
University of Tennessee, Knoxville, TN 37996

B. G. Cheon, J. S. Kang, K. Y. Kim
Korea University, Seoul 136-701, Korea

(E687 Collaboration)
(October 25, 1993)

An observation of an excited Λ_c^{+*} baryon decaying to $\Lambda_c^+ \pi^+ \pi^-$, with $\Lambda_c^+ \rightarrow p K^- \pi^+$ is presented. We reconstruct 39.7 ± 8.7 Λ_c^{+*} baryons with a mass of $340.4 \pm 0.6 \pm 0.3$ MeV/c² above the Λ_c^+ mass. The upper limit on the resonant branching ratio is $BR(\Lambda_c^{+*} \rightarrow \Sigma_c \pi^\pm) / BR(\Lambda_c^{+*} \rightarrow \Lambda_c^+ \pi^+ \pi^-) < 36\%$ at the 90% confidence level.

Charm spectroscopy has provided excellent tests of QCD quark models and Lattice Gauge calculations. The higher mass of the charm quark, relative to the strange quark, allows for more reliable results using QCD calculations. These tests have been limited mainly to $c\bar{c}$ meson spectroscopy. Data on charm baryon spectroscopy can be used to test further the agreement of various QCD mass predictions, in particular relativistic effects associated with the light quarks in the charm baryon.

Table I shows predictions for the masses of the $J^P = \frac{1}{2}^-$ and $\frac{3}{2}^-$ excited states of the Λ_c^+ and Σ_c baryons in a relativistic quark model with chromodynamics by Capstick and Isgur [1]. The Λ_c^{+*} baryon is forbidden to decay to $\Lambda_c^+ \pi^0$ since isospin is conserved in strong decays. If the Λ_c^{+*} has enough mass it can decay to $\Lambda_c^+ \pi^+ \pi^-$ (and $\Lambda_c^+ \pi^0 \pi^0$). Note that if the Σ_c^{+*} baryon has enough mass it can also decay to $\Lambda_c^+ \pi^+ \pi^-$ as well as $\Lambda_c^+ \pi^0$.

Evidence for an excited charm baryon state was first presented by ARGUS [2]. They reported a state decaying to $\Lambda_c^+ \pi^+ \pi^-$ with a mass of $2626.6 \pm 0.5 \pm 1.5$ MeV/c². They also report the branching fraction $BR(\Lambda_c^{+*} \rightarrow \Sigma_c \pi^\pm) / BR(\Lambda_c^{+*} \rightarrow \Lambda_c^+ \pi^+ \pi^-) = 0.51 \pm 0.23$. Although the observed signal could be due to either a Λ_c^* or a Σ_c^* baryon, it is almost certainly a Λ_c^* baryon, either the $\frac{1}{2}^-$ or/and the $\frac{3}{2}^-$ state because these are the two lowest lying states (see Table 1). These two states are predicted to have approximately the observed mass difference while the Σ_c^* states are predicted to have a much higher mass difference [1]. Furthermore, the Σ_c^* state should also be observable via its decay to $\Lambda_c^+ \pi^0$. Although a search for a Σ_c^{+*} state decaying to $\Lambda_c^+ \pi^0$ has not been reported, there is no evidence for the isospin related states decaying to $\Lambda_c^+ \pi^\pm$ [3]. We shall refer to the state first observed by ARGUS as the Λ_c^* or the $\Lambda_c(2625)$. Preliminary results from both CLEOII [4] and E687 [5] confirm the existence of the $\Lambda_c(2625)$. CLEOII reports 174 ± 21 events for $\Lambda_c^{+*} \rightarrow \Lambda_c^+ \pi^+ \pi^-$ with a mass difference of $342.1 \pm 0.4 \pm 0.5$ MeV/c² and 57 ± 15 events for $\Lambda_c^{+*} \rightarrow \Lambda_c^+ \pi^0 \pi^0$ with a mass difference of $340.7 \pm 0.9 \pm 1.4$ MeV/c².

This paper presents confirmation of a photoproduced $\Lambda_c(2625)$ decaying to $\Lambda_c^+ \pi^+ \pi^-$, where the Λ_c^+ is reconstructed via its decay to $p K^- \pi^+$. Throughout this paper, the charge conjugate state is implied when a decay mode of a specific charge is stated.

The data for this analysis were collected in 1990 and 1991 in the Fermilab wideband photoproduction experi-

ment E687. About 500 million triggers were recorded on tape during this period. The E687 detector is described in detail elsewhere [6].

A number of cuts were used to select $p K^- \pi^+$ combinations. Each of the decay secondaries must be reconstructed in both the microstrip and multiwire proportional chamber systems, and the two sets of track parameters must agree within measurement errors. Information from the Čerenkov counters is used to select protons, kaons, and pions.

The Λ_c^+ decays were reconstructed by a *candidate driven* method [6]. The three microstrip tracks of the $p K^- \pi^+$ combination forming the secondary vertex are required to extrapolate back to a single point with a confidence level greater than 1%. The candidate Λ_c^+ track must form a primary vertex with at least one other microstrip track with a confidence level greater than 1%. An important cut in isolating charm signals from non-charm background is the significance of detachment of the primary and secondary vertices. We use the variable ℓ/σ_ℓ , where ℓ is the signed three dimensional separation between the primary and secondary vertices, and σ_ℓ is the error on ℓ computed on an event by event basis taking into account the effects of multiple Coulomb scattering.

Figure 1(a) shows the $p K^- \pi^+$ invariant mass plot for $\ell/\sigma_\ell > 4$ and a momentum cut on the $p K^- \pi^+$ combination of $p_{\Lambda_c^+} > 50$ GeV/c. Monte Carlo studies show that $p K^- \pi^+$ combinations from Λ_c^+ baryons have a harder momentum spectrum than background $p K^- \pi^+$ combinations. The momentum cut keeps almost all the signal and improves the signal-to-noise so that a looser ℓ/σ_ℓ cut be used.

In the $p K^- \pi^+ \tilde{\pi}^+ \tilde{\pi}^-$ combination, the two *primary* ($\tilde{\pi}^+ \tilde{\pi}^-$) pions are required to be reconstructed in both the microstrip and the multiwire proportional chamber systems. In addition the microstrip tracks have to be part of the already reconstructed primary vertex. To reduce systematics we look at the mass difference between the $p K^- \pi^+ \tilde{\pi}^+ \tilde{\pi}^-$ combination and the $p K^- \pi^+$ combination: $\Delta M = M(p K^- \pi^+ \tilde{\pi}^+ \tilde{\pi}^-) - M(p K^- \pi^+)$. Note that for each event, all $\tilde{\pi}^+ \tilde{\pi}^-$ combinations are used with each $p K^- \pi^+$ combination.

Figure 1(b) shows the mass difference with a $\pm 2\sigma$ (± 19.2 MeV/c²) cut on the $p K^- \pi^+$ mass around 2.285 GeV/c². The mass difference distribution for background is shown by the dashed histogram. The background is from two sources: random $p K^- \pi^+$ combinations; and real Λ_c^+ baryons with random $\tilde{\pi}^+ \tilde{\pi}^-$ combinations. The random $p K^- \pi^+$ combination background is given by the (properly normalized) Λ_c^+ sidebands. A measure of the second background source is given by the $p K^- \pi^+ \tilde{\pi}^+ \tilde{\pi}^+$ and $p K^- \pi^+ \tilde{\pi}^- \tilde{\pi}^-$ combinations where (random) $p K^- \pi^+$ sideband contributions have been subtracted. A maximum likelihood fit, using a Gaussian signal and linear background, to the mass difference distribution yields 39.7 ± 8.7 events in the peak with a mass difference of $340.4 \pm 0.6 \pm 0.3$ MeV/c². The first error is

the statistical error and the second is the systematic uncertainty. The major contribution to the systematic uncertainty comes from fluctuations in the fitted mass difference for different data samples collected with various cuts, and for different fit functions and fit methods. Monte Carlo studies show that the shift in the mass difference due to acceptance is negligible ($0.06 \text{ MeV}/c^2$), and our (high statistics) data on the $M(D^0\pi^+) - M(D^0)$ mass difference agree with the world average value [7] to within $0.01 \text{ MeV}/c^2$.

The width of the peak is found to be $2.2 \pm 0.5 \text{ MeV}/c^2$ which is consistent with the value of $2.87 \pm 0.04 \text{ MeV}/c^2$ due to resolution alone, as determined in a Monte Carlo study.

Figure 1(c) shows the $pK^-\pi^+$ invariant mass when a cut is made in the ΔM mass difference of $\pm 2\sigma$ about $340.4 \text{ MeV}/c^2$. A fit to a Gaussian and linear background yields 47.9 ± 8.9 events. This gives further evidence that the peak in the mass difference is really associated with a Λ_c^+ .

The resonant decays $\Lambda_c^* \rightarrow \Sigma_c^0\pi^+$ and $\Lambda_c^* \rightarrow \Sigma_c^{++}\pi^-$ were investigated by plotting the $M(\Lambda_c^+\pi^+) - M(\Lambda_c^+)$ and $M(\Lambda_c^+\pi^-) - M(\Lambda_c^+)$ mass differences for a $\pm 2\sigma$ cut on the $pK^-\pi^+$ mass around $2.285 \text{ MeV}/c^2$ and a $\pm 2\sigma$ cut on the ΔM mass difference around $340.4 \text{ MeV}/c^2$. These mass difference plots are shown in Figure 2. In these plots, a fit function consisting of a quadratic background function plus two Gaussians was used. Due to the limited range of mass combinations constrained by the two mass cuts used, an actual $\Lambda_c^* \rightarrow \Sigma_c^{++}\pi^-$ event will also produce an entry in the $M(\Lambda_c^+\pi^-) - M(\Lambda_c^+)$ mass difference plot leading to a “false” peak at a mass of $340.4 - (M(\Sigma_c^{++}) - M(\Lambda_c^+)) \text{ MeV}/c^2$. Similarly a Σ_c^0 signal will give rise to a “false” peak in the $M(\Lambda_c^+\pi^+) - M(\Lambda_c^+)$ mass difference plot. For the fit, the widths of the real Σ_c in the mass difference plot were fixed at $1.77 \text{ MeV}/c^2$ and the widths of the “false” peaks were fixed at $2.84 \text{ MeV}/c^2$. These widths were obtained in a Monte Carlo study. The Σ_c peak mass difference values were also fixed in the fit to $167.8 \text{ MeV}/c^2$ and the false peak fixed to $172.6 \text{ MeV}/c^2$. Equal masses for the Σ_c^{++} and the Σ_c^0 have been assumed and an average of the world average values for these masses [7] and recent values obtained by CLEOII [8] were used. The background function used was obtained from a Monte Carlo study of non-resonant decays of the Λ_c^* .

The fit of the two histograms shown in Figure 2 was done in a combined maximum likelihood fit, so only three fit parameters were used in the fit of the two histograms: the number of Σ_c^{++} , the number of Σ_c^0 and the number of background events. The fit gave $4.1 \pm 3.2 \Sigma_c^{++}$ and $1.0 \pm 2.9 \Sigma_c^0$ events. A study of the Λ_c^* sidebands shows no evidence for (background) Σ_c^{++} or Σ_c^0 in our sample. Other fitting methods and fit functions gave fit values within one (statistical) sigma of these values. In particular a fit was tried where the number of Σ_c^{++} and the number of Σ_c^0 were constrained to be the same, this might be expected from isospin conservation if one ignores any

small differences in mass between the Σ_c^{++} and the Σ_c^0 . Due to the limited statistics and the fact that the resonant decay is consistent with zero, we prefer to quote an upper limit for the resonant decay. The actual limit was determined by studying the limits obtained using various cuts and fit methods and is corrected by 4.6% for the fact that only a $\pm 2\sigma$ region about the Λ_c^* mass was used in the analysis. The final 90% confidence level upper limit for the resonant decay is

$$\frac{BR(\Lambda_c^* \rightarrow \Sigma_c^{++}\pi^-) + BR(\Lambda_c^* \rightarrow \Sigma_c^0\pi^+)}{BR(\Lambda_c^* \rightarrow \Lambda_c^+\pi^+\pi^-)} < 36\%.$$

Theory predicts two Λ_c^* states ($1/2^-$ and $3/2^-$) split by spin-orbit interactions. While non-relativistic models predict relatively large spin-orbit splittings [9], relativistic models predict much smaller spin-orbit splittings [1]. We do not see any significant evidence of another peak in the mass difference distribution up to $450 \text{ MeV}/c^2$, whereas we might have expected to see both the $1/2^-$ and the $3/2^-$ states in this mass difference range. Three possible explanations are: the $1/2^-$ and $3/2^-$ splitting is too small to be resolved; the production of the two states is very different in our experiment; or the two states have very different branching ratios to $\Lambda_c^+\pi^+\pi^-$, (e.g. if one of the states has a low enough mass to be below the $\Lambda_c^+\pi^+\pi^-$ threshold).

It should be noted that there is some preliminary evidence from CLEO for a state decaying to $\Lambda_c^+\pi^+\pi^-$ with a mass difference of $308 \text{ MeV}/c^2$ [3]. However, the existence of this other state needs to be confirmed. Due to our limited statistics, the unknown relative production of the $340 \text{ MeV}/c^2$ and the $308 \text{ MeV}/c^2$ Λ_c^* states and the unknown branching ratios, we can neither confirm nor rule out the existence of this additional Λ_c^* state.

In conclusion, we confirm the existence of an excited charm baryon state at a mass of $340.4 \pm 0.6 \pm 0.3 \text{ MeV}/c^2$ above the Λ_c^+ mass. The upper limit on the resonant branching ratio to $\Sigma_c^{++}\pi^- + \Sigma_c^0\pi^+$ relative to $\Lambda_c^+\pi^+\pi^-$ is 36% at the 90% confidence level.

We wish to acknowledge the assistance of the staffs of the Fermi National Accelerator Laboratory, the INFN of Italy, and the physics departments of the collaborating institutions. This research was supported in part by the National Science Foundation, the U.S. Department of Energy, the Italian Istituto Nazionale di Fisica Nucleare and Ministero dell’Università e della Ricerca Scientifica e Tecnologica, and the Korean Science and Engineering Foundation.

* Now at Univ. of Maryland, College Park, MD 20742.

† Now at Univ. of Colorado, Boulder, CO 80309.

‡ Now at Dip. di Fisica Nucleare e Teorica dell’Università and INFN - Pavia, I-27100 Pavia, Italy.

[§] Now at Fermilab, Batavia, IL 60510.

^{**} Now at Yale Univ., New Haven, CT 06511.

- [1] S. Capstick and N. Isgur, Phys. Rev. D **34**, 2809 (1986).
- [2] ARGUS, H. Albrecht *et al.*, "Observation of a New Charmed Baryon", DESY Preprint DESY-93-101, July 1993.
- [3] M. Battle *et al.*, "Observation of a New Excited Charmed Baryon", contributed paper #303 submitted to Int. Symp. on Lepton and Photon Interactions, 10–15 August 1993, Ithaca, New York.
- [4] D. Acosta *et al.*, "Observation of the Excited Charm Baryon $\Lambda_c^{+\ast}$ in CLEOII", contributed paper #275 submitted to Int. Symp. on Lepton and Photon Interactions, 10–15 August 1993, Ithaca, New York.
- [5] J. Appel, plenary talk presented at Int. Symp. on Lepton and Photon Interactions, 10–15 August 1993, Ithaca, New York.
- [6] P. L. Frabetti *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **320**, 519 (1992).
- [7] Particle Data Group, K. Hikasa *et al.*, *Review of particle properties*, Phys. Rev. D **45**, 1 (1992).
- [8] B. Nemati, "Recent Charmed Baryon Results from CLEO", talk presented at the 7th Meeting of the APS Division of Particles and Fields, 10–14 November 1992, Fermilab, Batavia, Illinois.
- [9] L. A. Copley, N. Isgur and G. Karl, Phys. Rev. D **20**, 768 (1979).

FIG. 1. (a) $pK^-\pi^+$ invariant mass plot for $\ell/\sigma_\ell > 4$ and $p_{\Lambda_c^+} > 50$ GeV/c; (b) Using the data of (a), $\Delta M = M(pK^-\pi^+\bar{\pi}^+\bar{\pi}^-) - M(pK^-\pi^+)$ mass difference for $|M(pK^-\pi^+) - 2.285| < 19.2$ MeV/c² (circles) and for background (dashed histogram); (c) $pK^-\pi^+$ invariant mass plot as in (a) together with $|\Delta M - 0.3404| < 4.4$ MeV/c². The fits shown on the plots are to a Gaussian signal with a quadratic background (a) or a linear background function (b) and (c).

FIG. 2. (a) $M(\Lambda_c^+\pi^-) - M(\Lambda_c^+)$ mass difference; and (b) $M(\Lambda_c^+\pi^+) - M(\Lambda_c^+)$ mass difference. The fits shown are described in the text.

TABLE I. Mass predictions for the ground state and some excited states of the Λ_c^+ and Σ_c baryons taken from Ref. [1].

state	J^P	mass (MeV/c ²)	ΔM^a (MeV/c ²)
Λ_c	$\frac{1}{2}^+$	2265	
Λ_c^*	$\frac{3}{2}^-$	2630	365
Λ_c^*	$\frac{5}{2}^-$	2640	375
Σ_c^*	$\frac{3}{2}^-$	2765	500
Σ_c^*	$\frac{5}{2}^-$	2770	505

^aMass above the ground state of the Λ_c .




FIG. 1. (a) $pK^- \pi^+$ invariant mass plot for $l/\sigma_l > 4$ and $p_{A_c^+} > 50$ GeV/c ; (b) Using the data of (a), $\Delta M = M(pK^- \pi^+ \pi^+ \pi^-) - M(pK^- \pi^+)$ mass difference for $|M(pK^- \pi^+) - 2.285| < 19.2$ MeV/c^2 (circles) and for background (dashed histogram); (c) $pK^- \pi^+$ invariant mass plot as in (a) together with $|\Delta M - 0.3404| < 4.4$ MeV/c^2 . The fits shown on the plots are to a Gaussian signal with a quadratic background (a) or a linear background function (b) and (c).




FIG. 2. (a) $M(\Lambda_c^+ \pi^-) - M(\Lambda_c^+)$ mass difference; and (b) $M(\Lambda_c^+ \pi^+) - M(\Lambda_c^+)$ mass difference. The fits shown are described in the text.